AUTHOR INDEX

Ağargün, M.Y., 103
Agid, O., 360
Ahern, D.K., 179
Ahmad, F., 380
Akiskal, H.S., 13
Ali, S.O., 248
Alpers, D.H., 116
Alpert, J.E., 97
Alyanak, B., 216
Ampollini, P., 426
Andresen, B., 284
Andrews, G., 137
Apter, A., 35
Arduini, L., 446
Arnold, L.M., 83
Aronowitz, B., 4

Bachar, E., 360
Bagnato, M., 206
Bailey, E.D., 179
Bakim, B., 216
Baral, I., 216
Barsky, A.J., 179
Başoğlu, C., 103
Battaglia, M., 206
Bechtel, R.J., 326
Bell, A., 4
Benabarre, A., 339
Benazzi, F., 106
Berrios, G.E., 259
Berry, N.J., 24
Blundell, M.L., 373
Bogojevic, G., 28
Boiman, E., 167
Bonne, O., 360
Börner, I., 147
Borwn, P., 315
Bosc, M., 63
Bräunig, P., 147
Brown, E.S., 19
Burleson, J.A., 276

Cabras, P.L., 111
Canet, J., 172
Canetti, L., 360
Cantrell, P.J., 334
Carcamo, D., 326
Casati, J., 344
Cassano, B.G., 242
Çetin, M., 103
Chandragiri, S., 73
Cheng, D., 49
Cheng, Y., 57
Chignague, J.F., 369
Clark, A., 259
Cloninger, C.R., 426
Coccaro, E.F., 137
Cohen, L.J., 4
Colom, F., 339
Cox, C., 130

Dammen, T., 295
Daniels, B.A., 259
Delahousse, J., 369
Dell'Osso, L., 242
den Boer, J.A., 405
Denicoff, K.D., 248
De-Nour, A.K., 360
Di Bernardo, M., 111
Dolan, R., 315
Donati, D., 206
Donini, M., 206
Dowson, J.H., 42
Du, Z., 57
Duberstein, P.R., 130

Ebrinç, S., 103
Edmond, T., 49
Edvardsen, J., 416

Fama, J.M., 179
Fañanas, L., 380
Fava, M., 97
Favaretto, E., 123
Fink, M., 450
Fiorilli, M., 206
Fossati, A., 206
Francis, A., 73, 450
Frankenberg, F.R., 223
Franklin, D.L., 326
Friis, S., 295
Frye, M.A., 248
Fujimoto, K., 266
Fujita, H., 392
Fulton, M.K., 19

García, L., 172
Gariboldi, S., 426
Garno, J.L., 237
Gastó, C., 339
Gerhard, A., 77
Germain, A., 49
Ghaemi, S.N., 167
Giles, D.E., 130
Gillberg, C., 398
Gillberg, I.C., 398
Goldberg, J.F., 237
Goodwin, F.K., 167
Gothelf, D., 35
Gottschalk, L.A., 326
Gottwalz, E., 284
Green, R., 273
Grilo, C.M., 159, 163
Grounds, A.T., 42
Halstead, S.M., 70
Han, L., 57
Harrington, D.E., 326
Harris, V.L., 88
Hayashi, M., 197
Helenius, H.Y.M., 385
Hennen, J., 223
Hernandez-Avila, C.A., 276
Hiraga, M., 197
Höffler, J., 147
Hollander, E., 4
Hollifield, M., 49
Howard, R., 380
Hunter, E.E., 334

Igarashi, Y., 197
Ilonen, T., 385
Inoue, S., 392
Inoue, Y., 197
Ishige, N., 197
Ivarsson, T., 398
Iwasaki, Y., 266, 454
Izumoto, Y., 392

Joyce, P.R., 137

Karayiorgou, M., 373
Karlsson, H., 385
Kasai, S., 266
Keck, P.E., Jr., 83
Keller, M.B., 315
King, D.A., 130
Kingston, P., 4
Kiriike, N., 266, 454
Kirkby, K.C., 259
Koch, M., 73
Kocsis, J.H., 237
Kojima, M., 266
Kolko, D.J., 461
Koss, M., 49
Krakow, B., 49
Kranzler, H.R., 276
Krausz, M., 284
Kringlen, E., 416
Krüger, S., 147
Kwon, J., 4

Lalonde, B., 88
Landon, J.F., 334
Lasko, N.B., 24
Latas, M., 28
Lauerma, H., 385
Lavie, R., 35
Leinonen, K.-M., 385
Leon, A.C., 237
Lerer, B., 360

Leverich, G.S., 248
Levinson, D.M., 326
Liskow, B.I., 334
Loas, G., 369
Looper, K.J., 432
Lumley, M.A., 352
Lygren, S., 416

Macklin, M.L., 24
Madaro, D., 13
Maffei, C., 206
Maggini, C., 77, 426
Maguire, G.A., 326
Malur, C., 450
Mammen, O.K., 461
Mannucci, E., 111
Marchesi, C., 426
Marks, I., 273
Márquez, M., 172
Martin, G.A., 137
Martínez-Arán, A., 339
Masheb, R.M.., 159
Mataix-Cols, D., 273
Matsui, T., 266, 454
Matsunaga, H., 266, 454
Mavissakalian, M.R., 253
McElroy, L., 83
McElroy, S.L., 229
McGlashan, T.H., 163
Menzies, R.G., 259
Metzger, L.J., 24
Micheli, C., 13
Migone, P., 77
Mino, Y., 392
Miyata, A., 266
Modai, I., 289
Møller, P., 184
Moretti, S., 111
Moritz, S., 284
Mulder, R.I., 137
Murray, R.M., 380
Musetti, L., 13, 242

Nickel, E.J., 334
Nierenberg, A.A., 97
North, C.S., 116
Novella, L., 206
Nurnberg, H.G., 137

Offer, R., 35
Ohayon, M.M., 469
Øien, P.A., 416
Oidham, J.M., 137
Onstad, S., 416
Orr, S.P., 24
Ortiz, M., 172

Pallanti, S., 446
Paris, J., 432
Patten, S.B., 92
Pava, J., 97
Penick, E.C., 334
Perot, J.M., 369
Perugi, G., 13
Petrides, G., 73
Petty, F., 19
Phillips, K.A., 229, 315
Pilkonis, P.A., 461
Pini, S., 242
Pitman, R.K., 24
Poling, J., 276
Ponizovsky, A., 289
Portera, L., 237
Post, R.M., 248
Powell, B.J., 334
Primeau, F., 57
Prolo, F., 206
Pukrop, R., 438

Quilici, C., 13

Rankin, M.A., 97
Råstam, M., 398
Reinares, M., 339
Ricca, V., 111
Ritsner, M., 289
Rizvi, S., 73
Roberts, L.W., 303
Romano, A., 259
Rosenbaum, J.F., 97
Rosenthal, E., 57
Rossi, A., 446
Rotella, C.M., 111
Rounsaville, B.J., 276
Ruser, T., 223

Salokangas, R.K.R., 385
Salvador-Carulla, L., 172
Salvatore, P., 77
Şar, V., 216
Sarno, N., 242
Sass, H., 438
Schrader, R., 49
Seguí, J., 172
Selten, J.-P., 184
Semiz, Ü.B., 103
Shaley, A.Y., 360
Shapiro, C.M., 469
Shea, M.T., 315
Shimodera, S., 392
Shugar, G., 147
Shuriquie, N., 380
Sielky, K., 352
Skodol, A.E., 137
Skre, I., 416

Smith-Jackson, E.E., 248
Sobin, C., 373
Socci, C., 13
Sollinger, A.B., 248
Somoza, E., 137
Starcevic, B., 28
Steinmeyer, E.M., 438
Stratta, P., 446
Stubbs, J.H., 70
Sussams, P., 42

Taguchi, H., 197
Taiminen, T., 385
Talbot, N.L., 130
Tambs, K., 416
Tanaka, H., 454
Tandberg, d. 49
Taylor, J., 42
Tennen, H., 276
Toner, B.B., 344
Torgersen, S., 416
Torresani, S., 123
Trajkovic, G., 28
Trespalacios, H., 369
Trumpower, D., 303
Tuimala, P., 385
Tundo, A., 242
Tutkun, H., 216

Ujihara, H., 392
Uldall, K.K., 88

van den Bosch, R.J., 191
van Os, J., 380
Vieta, E., 339

Wallenius, E., 385
Wang, K., 57
Warner, T.D., 303
Warshaw, M.G., 315
Wentz, E., 398
Wiersma, D., 184
Wilberg, T., 295
Wilkeson, A., 19
Woodruff, P.W.R., 380
Wright, E.C., 97

Yamashima, M., 197
Yu, B., 344

Zamar, N., 253
Zanarini, M.C., 223
Zimmerman, C., 123
Zlotnick, C., 315
Zucchi, T., 111

SUBJECT INDEX

Absorption, in borderline personality disorder, 226
Acquired immune deficiency syndrome
  delirium in hospitalized patients with, 88-91
  morbidity in patients with, 9
Addiction Severity Index, higher scores associated with specific personality disorders, 277
Adjustment disorder, with depressed mood, in patients with binge eating disorder, 112
Adolescents
  ego defense mechanisms in healthy versus psychiatric inpatients, 35-41
  Israeli, impact of parental death versus separation on mental health, 360-368
  personality traits, relationship between TPQ and DSM-III-R, 426-431
Affectionless control rearing style, and drug-dependent patients, 123-129
Affective disorder
  association with personality disorder scores, 321
  and catatonia, 147
  see also Bipolar I disorder
Aggression, parent-to-child, and anger in mood and anxiety disorders, 461-468
Aggressive obsessions, in male OCD patients, 269
Agoraphobia
  in major depressive disorder, 97-102
  panic disorder patients, predictors of comorbid personality disorders in, 28-34
AIDS, *see* Acquired immune deficiency syndrome
Alcohol abuse, comorbiodity with social anxiety disorder, 407-408
Alcohol dependence
  in patients with binge eating disorder, 112
  and major depression, 94-95
  predictor of criminality, 276-284
Alcoholic patients, comorbid psychiatric diagnosis, 334-338
Alexithymia, gender and hemispheric functioning in, 352-359
Alproazolam, treatment, social anxiety disorder, 411
Alzheimer's Disease Assessment Scale-Cognitive Portion, scores for cognitive impairment compared with other instruments, 326-333
Amnesia, in borderline personality disorder, 226
AN, *see* Anorexia nervosa
ANAM, *see* Automated Neuropsychological Assessment Metric Battery
Anger
  and parent-to-child aggression in mood and anxiety disorders, 461-468
  in trichotillomania female patients, 347
Anhedonia, characterization of parasuicide, 369-372
Announcement, 401
Anorexia nervosa
  and depressive disorders, predictive factors, 401
  see also Eating disorders
  teenage-onset, depressive disorders in, 398-403
Antidepressants
  and cycle acceleration in bipolar I disorder, 17
  response of GAD and PD to, 253-258
  see also specific drugs
Antidepressants (*Continued*)
  treatment, to improve social functioning, 63-64
Antisocial personality disorder
  concordance rate in twin studies, 418 vcriteria for, 139
  impulsivity in, 433
  predictor of criminality, 277, 281
Antisocial personality trait
  association with worse long-term drinking outcome, 334-338
  TPQ dimensions in adolescents, 429
Anxiety
  psychological symptoms during adjustment in immigrants, 291, 292
  see also Social anxiety disorder
  variable in assessment of negative symptoms in schizophrenia, 191-196
Anxiety disorders
  anger and parent-to-child aggression in, 461-468
  association with personality disorders, trauma, and posttraumatic stress disorder, 315-325
  comorbidity
    in eating disorder patients, 454-460
    in major depression, 97-102
    with social anxiety disorder, 407-408
  insight and outcome in, 167-177
  prevalence in eating disorder patients, 456
ASI, *see* Addiction Severity Index
ASPD, *see* Antisocial personality disorder
Automated Neuropsychological Assessment Metric Battery, scores for cognitive impairment compared with other instruments, 326-333
Avoidant personality disorder
  concordance rate in twin studies, 418
  criteria for, 139
  in patients with body dysmorphic disorder, 4, 8, 231-233
Avoidant personality trait, TPQ dimensions in adolescents, 429
Axis I disorders, 296
  comorbidity in eating disorder patients, 454-460
Axis I and II disorders, comparison of Personality Diagnostic Questionnaire-4+ and Longitudinal, Expert, All Data diagnoses, 295-302

Barnes Akathisia Rating Scale, assessment of drug-induced akathisia, 70
BARS, *see* Barnes Akathisia Rating Scale
BDD, *see* Body dysmorphic disorder
BDI, *see* Beck Depression Inventory
BDII, *see* Bipolar II disorder
Beck Depression Inventory
  assessment of normal versus psychiatric inpatient adolescents, 35-41
  screening for binge eating disorder in obese outpatients, 111
  survey of seasonal changes in mood and emotion in Chinese college students, 57-62
BED, *see* Binge eating disorder
Behavior, seasonal change patterns of in Chinese college students, 57-62
Behavioral techniques, treatment, social anxiety disorder, 412

Benzodiazepines
response of GAD and PD to, 253-258
treatment
NMS, 75
social anxiety disorder, 411
BES, *see* Binge Eating Scale
Beta-blockers, treatment, social anxiety disorder, 411
Binge eating disorder
behavioral features, 159
body mass index in, 159-162
comparison of obese, nonobese, and bulimia nervosa, 160
mental disorders in patients with, 112
need for additional diagnostic criteria, 159-162
personality disorder criteria in, 163-166
screening obese outpatients for, 111-115
see also Bulimia nervosa; Eating disorders
Binge Eating Scale, screening for binge eating disorder in obese outpatients, 111
Bipolar disease, *see also* Manic-depressive
Bipolar disorder
associated with PTSD, 474
insight and outcome in, 167-177
neuroleptic exposure in outpatients, 248-252
psychotic
clinical characteristics, 242-247
psychopathological characteristics, 244
see also Manic symptoms
see also Mixed mania
with major depression, in teenage-onset anorexia nervosa, 398-403
Bipolar I disorder
clinical features in patients with different polarity from first episode, 16
demographic features of patients with different polarity of first episode, 15
depressive course
and fewer psychotic features, 16
and rapid cycling, 16
depressive onset and course of illness, 13-18
first-degree family history features in patients with different polarity from first episode, 16
insight and outcome in, 169
manic onset and course of illness, 13-18
mixed course and absence of rapid cycling, 17 vmixed onset and course of illness, 13-18
polarity of first episode and course of illness, 13-18
treatment with antidepressants, tendency for rapid cycling, 17
Bipolar II disorder
comorbidity with nonaffective disorders, 339-343
differential qualitative features in patients with and without comorbidity, 341
distinguished from borderline personality disorder, 106-110
insight and outcome in, 169
major depressive episode, comparison with unipolar disorder patients, 107
Bipolar mixed states, psychopathology of stable and unstable, historical view, 77-82
BITE, *see* Bulimic Investigation Test, Edinburgh
BMI, *see* Body mass index
BN, *see* Bulimia nervosa
Bodily sensation, amplification of as predictor of remission in DSM hypochondriasis, 179
Body dysmorphic disorder
axis I and axis II phenomena as distinct and dissociable, 10
cluster C personality disorders in, 9
comorbid personality impairment in, 4-12
form of obsessive-compulsive disorder, 234
personality disorders in patients with, 229-236
similarity to OCD, 4-5
Body image, in trichotillomania female patients, 348. See also Body dysmorphic disorder
Body mass index
comparison in patients with and without binge eating disorder, 112, 113
in binge eating disorder, 159-162
Bonn Scale for the Assessment of Basic Symptoms, measurement of schizophrenia basic symptoms, 284-285
Borderline personality disorder
associated with
BDD, 4, 8
with PTSD, 315-325
concordance rate in twin studies, 418
criteria for, 139
dissociation in as causal factor for psychotic phenomena, 42-48
dissociative experiences of patients, 223-227
distinguished from bipolar II disorder, 106-110
in female obsessive-compulsive disorder patients, 266-272
gender-related in OCD, 270
impulsivity in, 433
predictive power of self-reported past psychotic phenomena for DSM-III-R diagnosis, 47
relationship to childhood sexual abuse, 131
wide range of dissociative experiences in, 224-226
Borderline personality trait, TPQ dimensions in adolescents, 429
BPD, *see* Borderline personality disorder
BPRS, *see* Brief Psychiatric Rating Scale
Brain, interhemispheric transfer deficit in alexithymia, 355-356
Brief Psychiatric Rating Scale, assessment of
basic symptoms of schizophrenia, 285-286
psychopathology in psychotic biopolar disorder, 242-247
Brief Symptom Inventory, assessment of effects of parental loss in adolescents, 362
Brofaromine, treatment, social anxiety disorder, 411
Bronazepam, treatment, social anxiety disorder, 411
BSABS, *see* Bonn Scale for the Assessment of Basic Symptoms
BSI, *see* Brief Symptom Inventory
Bulimia nervosa, diagnostic criteria, 159-162. *See also* Binge eating disorder
Bulimic Investigation Test, Edinburgh, screening for binge eating disorder in obese outpatients, 111

Caffeine, induction of panic attacks in patients with social anxiety disorder, 408
Camberwell Family Interview, association between distress and expressed emotion in family members of schizophrenia patients, 392-397
Carbamazepine
exposure to in bipolar disorder outpatients, 249-250
response to in mixed mania, 86

Carbon dioxide, induction of panic attacks, 408
CAS, *see* Community Adaptation Schedule
Case reports
  pseudoakathisia, 70-72
  therapeutic effects of delirium in psychosis, 450-452
Catatonia
  association with neuroleptic malignant syndrome, 73-75
  characteristics, 147
  comparison of motor and behavioral symptoms, 148
  GABA-ergic systems in, 75
  score frequency and severity range for each item on CRS, 152
Catatonia Rating Scale
  advantages, 152-153
  diagnostic threshold, 149-150
  instrument, 154-157
  item selection, 148-149
  severity rating, 149
  time, 150
Catatonia Rating Scale I, psychometric properties, 147-158
Cerebrum, structural abnormalities, association with dermatoglyphic ridge counts in schizophrenia, 380-384
CFI, *see* Camberwell Family Interview
CGI, *see* Clinical Global Impression
Childhood abuse, in dissociative disorders, psychiatric outpatients in Turkey, 216-222
Childhood sexual abuse
  association with anxiety disorder, 317
  definition, 131
  personality traits of women with history of, 130-136
Chlorpromazine, effects on basic symptoms of schizophrenia, 284-288
CIDI, *see* Composite International Diagnostic Interview
Circumplex models, personality disorders, 438-445
Civilian trauma
  psychiatric sequelae, 19-23
  psychopathology following, 20
Clinical Global Impression, assessment of
  insight and outcome in bipolar, unipolar, and anxiety disorders, 167
  suicidal behavior in patients with panic disorder, 103
Clinical screening instrument, development of for personality disorders, 137-146
Clinician Rating of Drinking Scale, assessment of comorbid psychiatric diagnosis and long-term drinking outcome, 335
Clomipramine, treatment, social anxiety disorder, 409
Clonazepam, treatment, social anxiety disorder, 411
Cloninger's neruobiologic learning model, support for in classification of personality variants, 426-431
Cluster B personality disorders, dimensions underlying, 432-437
Cocaine dependence, interaction with antisocial personality disorder, 281
Cognitive impairment, computerized measurement, 326-333
Cognitive-behavioral therapy, treatment social anxiety disorder, 412
Combat-related trauma, personality disorders associated with, 322-323
Community Adaptation Schedule, assessment of social functioning, 64
Comorbidity
  alcohol abuse and social anxiety disorder, 407-408
  alcoholism and psychistric diagnosis, 334-338
  anxiety disorder
    and eating disorder, 454-460
    and major depression, 97-102
    and social anxiety disorder, 407-408
  axis I disorders in eating disorder patients, 454-460
  bipolar disorder and PTSD, 474
  bipolar II disorder , 341
    and nonaffective disorders, 339-343
  body dysmorphic disorder and personality impairment, 4-12
  depression and anxiety disorders, 97-102
  depressive disorders and anorexia nervosa, 398-403
  dissociative disorders, 219
  dysthymia and PTSD, 474
  effect on treatment strategies for social anxiety disorder, 412-413
  major depression
    and eating disorder, 454-460
    and PTSD, 474
    and social anxiety disorder, 407-408
  major depressive disorder and PTSD, 474
  mood disorder and eating disorder, 454-460
  obsessive-compulsive disorder
    and body dysmorphic disorder, 231-233
    and gender identity disorder, 273-275
    and major depressive disorder, 97-102
    and panic disorder, 174
  panic disorder, 174
    and binge eating disorder, 112
  panic disorder/agoraphobia, 28-34
  paranoid personality disorder and BDD, 4, 231-233
  passive-aggressive personality disorder and BDD, 4, 8
  personality disorders and
    panic disorder/agoraphobia, 28-34
    BDD, 229-236
  posttraumatic stress disorder and other mental disorders, 475
  psychiatric diagnosis and alcoholism, 334-338
  psychoactive substance use disorder and eating disorder, 454-460
  schizotypal personality disorder and male OCD, 266-272
  substance abuse and social anxiety disorder, 407-408
Composite International Diagnostic Interview, risk factors for major depression, 92-96
Computer treatment, in obsessive-compulsive disorder, 259-265
Computerized measurement, psychiatric states, 326-333
Control in trichotillomania female patients, 348
Controlled Oral Word Association Test, scores for cognitive impairment compared with other instruments, 326-333
Cortisol, plasma and cerebrospinal fluid, elevated levels in mixed mania, 85
Criminality
  predictors of, personality and substance use disorders, 276-284
  risk factors for in personality and substance abuse disorder patients, 280
CRS, *see* Catatonia Rating Scale
Culture, differences in cluster B personality disorders, 435

Dam collapse, MDD and GAD following, 21
DAPI, *see* Dimensional Assessment of Personality Impairment
DDIS, *see* Dissociative Disorders Interview Schedule
Defense mechanisms
  correlations between and BDI, STAI, and MAI scores, 39
  in healthy versus psychiatric inpatient adolescents, 35-41
  specific, related to specific emotions, 39
Delerium
  in hospitalized AIDS patients, 88-91
  therapeutic in psychosis, 450-453
Delusional disorder, in patients with binge eating disorder, 112
Delusions, self-reported, correlations with DSM-III-R personality disorders, 45
Dementia praecox, differentiation from manic-depressive illness, historical view, 77-82
Dependent personality disorder
  associated with BDD, 4, 8
  concordance rate in twin studies, 418
  criteria for, 139
  in female obsessive-compulsive disorder patients, 266-272
  in patients with body dysmorphic disorder, 231-233
Dependent personality trait, TPQ dimensions in adolescents, 429
Depersonalization
  comorbidity with simple phobia in panic disorder, 174-175
  in borderline personality disorder, 226
  in panic disorder, 172-178
  symptoms associated with, 176
Depression
  anhedonia as in parasuicides, 369-372
  assessment of social functioning in, 63-69
  associated with BDD, 4, 8
  characteristics in psychotic bipolar disorder, 242-247
  following civilian trauma, 21
  and loss of parent during adolescence, 360-368
  major
    age- and sex-adjusted exposure frequencies, 95
    comorbid anxiety disorders in, 97-102
    demographic characteristics of clinical subjects, 93-94
    differences in psychological symptoms in patients with comorbid anxiety disorders, 99, 100
    live events, traumatic experiences, and social support associated with, 93
    mood disorder risk factors, 94
    specificity of epidemiological correlates, 92-96
  most common onset in bipolar I disorder, 14, 15
  psychological symptoms during adjustment in immigrants, 291, 292
  recovery from and social functioning, 63-64
  see also Bipolar I disorder; Major depressive disorder
  triggered by anorexia nervosa, 398-403
  use of SASS as outcome measure in, 66-68
  variable in assessment of negative symptoms in schizophrenia, 191-196
Depressive disorders
  and anorexia nervosa, predictive factors, 401
  associated with PTSD, 474
  in teenage-onset anorexia nervosa, 398-403
Depressive temperament, and hyperthymic, associated with depressive onset in bipolar I disorder, 15
Dermatoglyphic ridge counts, association with cerebral structural abnormalities in schizophrenia, 380-384
DES, *see* Dissociative Experiences Scale
Dexamethasone suppression test, nonsuppression in cases of mixed mania, 85
Diagnosis, social anxiety disorder, 406-408
Diagnostic criteria, need for additional in BED, 159-162
Diagnostic Interview for DSM-IV Personality Disorders, validity of criteria for personality disorders in BED, 163-166
Diagnostic and Statistical Manual-IV, criteria, use to distinguish borderline personality disorder from bipolar II disorder, 108
Dimensional Assessment of Personality Impairment
  methods, evaluation of personality impairment in BDD patients, 4-12
  reliability and validity in testing, 7-8
DIPD-IV, *see* Diagnostic Interview for DSM-IV Personality Disorders
Dissociative Disorders Interview Schedule, diagnosis of dissociative disorders in psychiatric outpatients in Turkey, 217
Dissociative disorders
  comorbidity and subdiagnoses, DDIS, 219
  frequency in psychiatric outpatients in Turkey, 216-222
  main symptom clusters, DDIS, 218
  mental health characteristics, DDIS, 219
Dissociative experiences, in borderline personality disorder patients, 223-227
Dissociative Experiences Scale
  borderline personality patients, 223-227
  screening for chronic dissociative disorders in psychiatric outpatients in Turkey, 217
Distress
  changes in symptoms of during adjustment of immigrants, 289-294
  family, in relative of schizophrenia patients, 392-397
  immigrants during adjustment, changes in frequency of psychological symptoms, 292
Dopamine transporter, striatal presynaptic, dysfunction in social anxiety disorder, 405-415
Dopaminergic function, in social anxiety disorder, 408
Dopaminergic systems, dysfunction in social anxiety disorder patients, 409
Drive-mood system, manic-depression, Hamburg school, 78-79
DRP, *see* Duration of prodrome
Drug abuse, measurement of cognitive impairment and psychiatric states by computer, 326-333. *See also specific drugs*
Drug Attitude Inventory, subjective response to neuroleptics in schizophrenics, 446-449
Drug dependence
  affectionless control rearing style in, 123-129 vin panic disorder, 174
  parental representations in, 123-129
  predictor of criminality, 276-284
  see also specific drugs
DSM-IV criteria, *see* Diagnostic and Statistical Manual-IV
DST, *see* Dexamethasone suppression test
DUP, *see* Duration of untreated psychosis
Duration of prodrome, in first-episode schizophrenia, 186-187

Duration
of untreated prognosis, relationship between duration of and psychosis in schizophrenia, 184-190
of untreated psychosis
first-episode schizophrenia, temporal construction, 186
and symptom content, first-episode schizophrenia, 188-189
Dynamic psychotherapy, treatment, social anxiety disorder, 412
Dysphoria, manic-depressive, Berner's conception, 79-81
Dysthymia
associated with PTSD, 474
in panic disorder, 174
in patients with binge eating disorder, 112

Eating Disorder Examination—Questionnaire, diagnosis of binge eating disorders, 160
Eating disorders
comorbidity of axis I disorders, 454-460
in panic disorder, 174
see also Anorexia nervosa; Binge eating disorder; Bulimia nervosa
Eccentric personality disorder, concordance rate in twin studies, 418
ECT, *see* Electroconvulsive therapy
EDE-Q, *see* Eating Disorder Examination—Questionnaire
EDS, *see* Ego Defense Scale
Efficiency trials, controlled, generalized anxiety disorder versus panic disorder, 253-258
Ego defense mechanisms, *see* Defense mechanisms
Ego Defense Scale, assessment of defense mechanisms in normal and psychiatric inpatient adolescents, 35-41
Electroconvulsive therapy, induction of incidental deliurium in psychosis, 450-453
Embarrassment, in trichotillomania female patients, 346
EMDR, *see* Eye movement desensitization and reprocessing
Emotional personality disorder, concordance rate in twin studies, 418
Emotions
psychoevolutionary theory of, 35
see also Alexithymia; Expressed emotion
specific, related to specific defense mechanisms, 39
Epidemiological correlates, specificity of in major depression, 92-96
Epidemiology, social anxiety disorder, 406
Expressed emotion, in relatives of schizophrenia patients, 392-397
Extrapyramidal symptoms, and neuroleptic treatment of basic symptoms of schizophrenia, 284-288
Extraversion
low in patients with body dysmorphic disorder, 232-233
scores for in women with history of childhood sexual abuse, 132
Eye movement desensitization and reprocessing, therapy for combat-related PTSD, 24-27

FBF, *see* Franfurter Beschwerde-Fragebogen
FCQ, *see* Frankfurt Complaint Questionnaire
Fear
response, in social anxiety disorder, 408
in trichotillomania female patients, 347
Fearful personality disorder, concordance rate in twin studies, 418
Finger Tactile Localization Test, assessment of hemispheric functioning and gender in alexithymia, 352-359
Five-Minute Speech Sample, association between distress and expressed emotion in family members of schizophrenia patients, 392-397
Fluoxetine
assessing efficacy of in improving social function in patients with depression, 63-69
compared with reboxetine in improving social functioning, 66
Fluvoxamine, response to by patients with body dysmorphic disorder, 233
FMSS, *see* Five-Minute Speech Sample
Frankfurt Complaint Questionnaire, assessment of basic symptoms of schizophrenia, 284-285
Frankfurter Beschwerde-Fragebogen, assessment of subjective experience of psychosis in schizophrenia, 446-449
Frustration, in trichotillomania female patients, 347

GABA, *see* Gamma aminobutyric acid
Gabapentin, treatment, social anxiety disorder, 412
GAD, *see* Generalized anxiety disorder
GAF, *see* Global Assessment of Functioning
GAFS, *see* Global Assessment of Functioning Scale
Gamma-aminobutyric acid systems, and pathophysiology of catatonia, 75
Gender
and criminality in personality disorder patients, 278-279
differences in
cluster B personality disorders, 434-435
patients with obsessive-compulsive disorder, 266-272
and hemispheric functioning, in alexithymia, 352-359
influence over outcome in mixed mania, 85-86
OCD in male schizotypal personality disorder, and PTSD, 472
potential confounding effect, analysis of patterns of covariation of DSM-IV personality disorders in mixed psychiatric population, 206, 212
and prevalence of binge eating disorder in obese outpatients, 113
prevalence of PTSD by, 472
role in mixed mania, 83-87
role in seasonal affective disorder, study of Chinese college students, 57-62
see also Gender identity disorder; Women
and social functioning impairment in OCD, 269
Gender identity disorder
case series, 274
comorbidity with obsessive-compulsive disorder, 273-275
remission in, 273-275
treatment methods, 273-274
General Health Questionnaire, association between distress and expressed emotion in family members of schizophrenia patients, 392-397
General Well Being Schedule, assessment of effects of parental loss in adolescents, 362
Generalized anxiety disorder
characteristics of patients, 255, 256
following civilian trauma, 21
in major depressive disorder, 97-102
in patients with binge eating disorder, 112

Generalized anxiety disorder (*Continued*)
versus panic disorder, in controlled efficiency trials, 253-258
in Vietnam War veterans, 22
GHQ, *see* General Health Questionnaire
GID, *see* Gender identity disorder
Global Assessment of Functioning Scale
assessment of
functioning in OCD, 267
insight and outcome in bipolar, unipolar, and anxiety disorders, 167
use in distinguishing borderline personality disorder from bipolar II disorder, 107
Global Seasonality Score, assessment of seasonal changes in mood and emotion in Chinese college students, 57-62
Gottschalk CI Scale, revalidation, 330-331
Grandiosity
and duration of untreated psychosis in first-episode schizophrenia, 187
in mixed versus pure mania, 239
GSS, *see* Global Seasonality Score
Guilt, in trichotillomania female patients, 347
GWB, *see* General Well Being Schedule

Hair, *see* Trichotillomania
Hallucinations
and borderline personality disorders, 43
in PTSD, 473
self-reported, correlations with DSM-III-R personality disorders, 45
Haloperidol, akathisia as side effect, 70
Hamburg school, mixed states and mixed pictures, manic-depressive, 78-79
Hamilton Anxiety and Depression Rating Scale, assessment of depersonalization in panic disorder, 172-173
Hamilton Depression Rating Scale, diagnosis of mixed mania, 84
Hamilton Rating Scale for Depression assessment of comorbid anxiety disorders in MDD, 98
Harm avoidance, TPQ dimensions in adolescents, 426-431
HCI, *see* Human-computer interactions
Hemispheric functioning, and gender in alexithymia, 352-359
Heritability, personality disorders, 415-425
History, psychopathology of stable and unstable mixed states, 77-82
Histrionic personality disorder
associated with BDD, 4, 87
concordance rate in twin studies, 418
criteria for, 139
impulsivity in, 433
Histrionic personality trait, TPQ dimensions in adolescents, 429
Hostility, psychological symptoms during adjustment in immigrants, 291, 292
Human-computer interactions, as process measure in psychotherapy patients, 259-265
Humiliation, in trichotillomania female patients, 347-348
Hyperthymic temperament, associated with manic course in bipolar I disorder, 15
Hypochondriasis
disorder or perception and cognition, 181
DSM, predictors of remission in, 179-183
logistic regression model of status, 181
Hypothyroidism, associated with rapid cycling in mixed mania, 85

IBS, *see* Irritable bowel syndrome
Immigrants, changes in psychological symptoms during adjustment, 289-294
Impulse-control disorder, *see* Trichotillomania
Impulsive traits, severity in cluster B personality disorders, 432-437
Impulsivity, and outcome in cluster B personality disorders, 433
Incest, parental, and personality traits of women who experienced as child, 130-136
Information processing, in first-episode schizophrenia, 385-391
Insight, and outcome in bipolar, unipolar, and anxiety disorders, 167-177
Insomnia
assessment of in sexual assault survivors, 51
following sexual assault, 49-56
in PTSD, 473
Insulin coma therapy, therapeutic effects in psychosis, 450-452
Irritability, in mixed versus pure mania, 239
Irritable bowel syndrome
associated with psychiatric diagnosis, 119
criteria for, 118
in psychiatric patients, 116-122
symptoms in psychiatric patients, 119
Isolation, in trichotillomania female patients, 346-347

LEAD, *see* Longitudinal, Expert, All Data
Life events, associated with major depression, 93
Life Experience Scale, assessment of personality traits of women with history of childhood sexual abuse, 132
Life Style Index, assessment of defense mechanisms in normal and psychiatric inpatient adolescents, 35-41
Lithium
exposure to in bipolar disorder outpatients, 249-250
treatment, response to in mixed mania, 86
Longitudinal, Expert, All Data, compared with PDQ-4+ in diagnosis of axis I and II disorders, 295-302
LSI, *see* Life Style Index

MADRS, *see* Montgomery and Asberg Depression Rating Scale
Magnetic resonance imaging, assessment of dermatoglyphic ridge counts and cerebral structural abnormalities in schizophrenia, 380-384
MAI, *see* Multidimensional Anger Inventory
Major depression
comorbidity in eating disorder patients, 454-460
in panic disorder, 174
in patients with binge eating disorder, 112
Major depression, *see* Depression, major
Major depressive disorder
assessment of defense mechanisms in adolescents with, 35-41
associated with PTSD, 474
association with personality disorder scores, 321
comorbidity with social anxiety disorder, 407-408
following civilian trauma, 21
in Vietnam War veterans, 22
Major depressive disorder, *see also* Depression, major

Mania
characteristics in psychotic bipolar disorder, 242-247
mixed
characteristics in psychotic bipolar disorder, 242-247
difficulty in recognizing, 240-241
role of gender, 83-87
versus pure, manic features during, 239
versus pure, qualitative differences in manic symptoms during, 237-241
pure and mixed, demographic, clinical, and treatment characteristics, 239
Manic depressive, mixed states in Kraepelin and Weygandt, 77-78
Manic symptoms, qualitative differences in during mixed versus pure mania, 237-241
MAOIs, *see* Monoamine oxidase inhibitors Marks and Mathews Fears and Phobias Scale, assessment of depersonalization in panic disorder, 172-173
MDD, *see* Major depressive disorder
Medical Outcomes Study, assessment of social functioning, 64
Men, alexithymia and hemispheric functioning in, 356-357
Menstrual cycle, relationship to suicidal behavior in patients with panic disorder, 103-105
Mental disorders, associated with PTSD, 474
Mental health measures
adolescents experiencing parental loss, 364
adolescents experiencing parental separation, 365
Minnesota Multiphasic Personality Inventory, assessment of comorbid psychiatric diagnosis and long-term drinking outcome, 335
Mixed mania
biology of, 85
criteria for defining, 83, 84, 85
predictors for outcome, 85-86
response to carbamazepine treatment, 86
response to lithium treatment, 86
role of gender, 83-87
Mixed pictures, Hamburg school, 78-79
Mixed states
Hamburg school, 78-79
manic-depressive, in Kraepelin and Weygandt, 77-78
stable, Berner, 81
unstable, Berner, 79-81
MMPI, *see* Minnesota Multiphasic Personality Inventory
Moclobemide, treatment, social anxiety disorder, 411
Monoamine oxidase inhibitors, treatment, social anxiety disorder, 409-411
Montgomery and Asberg Depression Rating Scale
assessment of anhedonia in parasuicides, 369
use in distinguishing borderline personality disorder from bipolar II disorder, 107
Mood, seasonal change patterns of in Chinese college students, 57-62. *See also* Seasonal affective disorder
Mood disorder
anger and parent-to-child aggression in, 461-468
comorbidity in eating disorder patients, 454-460
see also Borderline personality disorder
MOS, *see* Medical Outcomes Study
Motor vehicle accidents, MDD and substance abuse following, 21
Movement disorders, *see* Pseudoakathisia
Multidimensional Anger Inventory, assessment of normal and psychiatric inpatient adolescents, 35-41

Narcissistic personality disorder
associated with BDD, 4, 8
concordance rate in twin studies, 418
criteria for, 139
impulsivity in, 433
predictor of criminality, 277
TPQ dimensions in adolescents, 429
NART, *see* National Adult Reading Test
National Adult Reading Test, assessment of computer-aided treatment of OCD, 259-265
Nefazodone, treatment, social anxiety disorder, 412
Negative emotions, and defense mechanisms in adolescents, 35
Negative symptoms, discrepancy between examiner rating and self-rating, 191-196
Neglect, childhood, in dissociative disorders, psychiatric outpatients in Turkey, 220
NEO-FFI, *see* Neuroticism, Extraversion, Openness Five Factor Index
Neuroendocrine function, in social anxiety disorder, 408
Neuroimaging, of social anxiety disorder patients, 409
Neuroleptic malignant syndrome
association with catatonia, 73-75
severity of and number of catatonic signs, 73-75
Neuroleptics
effect of dosage on basic symptoms of schizophrenia, 284-288
exposure to in bipolar disorder outpatients, 248-252
prospective use in bipolar patients, 250
subjective experience and response to in schizophrenia, 446-449
Neuroticism, Extraversion, Openness Five Factor Index
administered to BDD patients, 5
assessment of
personality disorders in patients with body dysmorphic disorder, 230-231
personality traits of women with history of childhood sexual abuse, 130
Neuroticism, high in patients with body dysmorphic disorder, 231-233
Nightmares, in PTSD, 473
NMS, *see* Neuroleptic malignant syndrome
Noradrenergic function, in social anxiety disorder, 408
Novelty seeking, TPQ dimensions in adolescents, 426-431

Obesity, screening for binge eating disorder in, 111-115
Obsessive-compulsive disorder
assessment of defense mechanisms in adolescents with, 35-41
clinical characteristics
in adults without childhood onset, 375
in children, 374
comorbidity
in eating disorder patients, 454-460
with gender identity disorder, 273-275
computer treatment process, 259-265
concordance rate in twin studies, 418
criteria for, 139
early- and late-onset, phenotypic differences, 373-379
gender differences in patients with, 266-272

Obsessive-compulsive disorder (*Continued*)
in major depressive disorder, 97-102
in panic disorder, 174
in patients with body dysmorphic disorder, 231-233
similarity to BDD, 4-5
Obsessive-compulsive personality trait, TPQ dimensions in adolescents, 429-430
Obsessiveness, psychological symptom during adjustment in immigrants, 291, 292
Occupational functioning, gender-related impairment in OCD, 269
OCD, *see* Obsessive-compulsive disoder
Openness to experience, score for in women with history of childhood sexual abuse, 132-133
Opioid dependence, interaction with personality disorders, 281

Padus Inventory, assessment of computer-aided treatment of OCD, 259-265
Pain, in trichotillomania female patients, 347-348
Panic attacks, substances that induce, 408-409
Panic disorder
characteristics of patients, 255-256
clinical characteristics, 174
comorbidity, 174
comparison of patients with and without personality disorder, 30
depersonalization in, 172-178
implication of temporal lobe in pathogenesis of, 176
in major depressive disorder, 97-102
in patients with binge eating disorder, 112
predictors of
DSM-IV Cluster A personality disorder in, 31
DSM-IV Cluster B personality disorder in, 31
DSM-IV Cluster C personality disorder in, 31
personality disorder in, 30
premenstrual exacerbation and suicidal behavior in patients with, 103-105
Spearman correlations between test scores, 32
versus generalized anxiety disorder, in controlled efficiency trials, 253-258
Panic disorder/agoraphobia
patients, predictors of comorbid personality disorders in, 28-34
personality disorders associated with high levels of parental protection, 28-34
Panic-Associated Symptom Scale, assessment of depersonalization in panic disorder, 172-173
Paranoid ideation, psychological symptoms during adjustment in immigrants, 291, 292
Paranoid personality disorder
associated with BDD, 4
criteria for, 139
in patients with body dysmorphic disorder, 231-233
predictor of criminality, 277
symptoms, self-reported, correlations with DSM-III-R personality disorders, 46
Paranoid personality trait, TPQ dimensions in adolescents, 427-428
Parasuicide, and anhedonia, 369-372
Parental Bonding Instrument
analysis of personality disorder in PDAG, 28-34
Parental Bonding Instrument (*Continued*)
assessment of effects of parental loss in adolescents, 362
assessment of parental representations in drug-dependent patients, 123-129
Parental care, lower levels in depressed patients with comorbid personality disorders, 28
Parental protection
correlation with DSM-IV personality disorder clusters, 31
high levels of associated with personality disorders in PDAG, 28
and personality disorder in PDAG patients, 32-33
Parenting, quality of in drug-dependent patients, 123-129
Parents
Aggression and anger in mood and anxiety disorders, 461-468
impact of death versus separation on mental health of Israeli adolescents, 360-368
parental representations in drug-dependent patients, 123-129
Parent-to-child aggression, and anger in mood and anxiety disorders, 461-468
Paronoid personality disorder, concordance rate in twin studies, 418
Paroxetine, treatment, social anxiety disorder, 410
PASS, *see* Panic-Associated Symptom Scale
Passive-aggressive personality disorder
associated with BDD, 4, 8
concordance rate in twin studies, 418
criteria for, 139
TPQ dimensions in adolescents, 430
PBI, *see* Parental Bonding Instrument
PCA, *see* Principal component analysis
PDAG, *see* Panic disorder/agoraphobia
PDI, *see* Psychiatric Diagnostic Interview
PDQ-4+, *see* Personality Diagnostic Questionnaire-4+
Pentagastrin, induction of panic attacks in social anxiety disorder patients, 409
Perceived Immigration-Related Stressors Scale, assessment of psychological symptoms during adjustment in immigrants, 289-290
Perceived Social Support Family/Friends Scale, assessment of effects of parental loss in adolescents, 362
Performance, effective, impaired in first-episode schizophrenia, 385-391
Persistence, TPQ dimensions in adolescents, 426-431
Personal and Social Network Adjustment Scale, assessment of social functioning, 64
Personality
comorbid impairment, in BDD, 4-12
study of defense mechanisms in, 35
Personality Diagnostic Questionnaire-4+, compared with LEAD in diagnosis of axis I and II disorders, 295-302
Personality disorder
according to gender in obsessive-compulsive disorder, 266-272
borderline, distinguished from bipolar II disorder, 106-110
characteristics of best discriminative criteria, 141, 144
demographic data, 173
DSM-IV, correlation with parental protection, 31
see also specific types

Personality Disorder Examination, assessment in
traumatic personality, 316-317
personality disorders, 137
Personality disorders
adverse effect on treatment response, 10
associated with self-reported past psychotic phenomena, 42-48
association
of scores with diagnosis of MDD, 321
with anxiety disorders, 315-325
in BED, internal consistency, MICs, and ICMICs for, 165
big-five model, 438-440
categorical-typological approach, 438
circumplex models, 438-445
cluster B
dimensions underlying, 432-437
factors differentiating, 433-435
cluster C, in BDD, 9
comorbid, in panic disorder/agoraphobia patients, 28-34
criteria for, 318-321
development of clinical screening instrument for identifying, 137-146
dimensional approach, 438-440
discriminative criteria for, 139
DSM-III-R, correlations
with self-reported hallucinations, 45
with self-reported delusions, 45
DSM-IV
Axis II, validity of criteria for in BED, 163-166
latent dimensions not described by, 214
patterns of covariation in mixed psychiatric sample, 206-215
heritability of, 415-425
model testing in twin studies, 420-422
in patients with body dysmorphic disorder, 229-236
predictors of criminality, 276-284
rates of in cases of trauma, 318
twin study, 416-425
Personality traits
adolescents, relationship between TPQ and DSM-III-R, 426-431
of women with history of childhood sexual abuse, 130-136
Phenelzine, treatment, social anxiety disorder, 411
Phobias, in panic disorder, 174
Physical Anhedonia Scale, assessment of anhedonia in parasuicides, 369
Physical force, in childhood sexual abuse, and personality traits of women who experienced as child, 131
PI, *see* Padus Inventory
PIRSS, *see* Perceived Immigration-Related Stressors Scale
Planning, impaired in first-episode schizophrenia, 385-391
Polysomnography, assessment algorithm for in sleep disorders following sexual assault, 52
Posttraumatic stress disorder
as result of civilian trauma, 19-23
association with anxiety disorders, 315-325
axis II features in, 315-325
combat-related, EMDR therapy for, 24-27
comorbidity with other mental disorders, 475
prevalence, 475
by gender, 472
Posttraumatic stress disorder (*Continued*)
sexual-assault survivors, sleep breathing and sleep movement disorders in, 49-56
sleep disorders in, 475-476
and psychiatric disorders associated with, 469-478
and social network, 474
in Vietnam War veterans, 22
Premenstrual exacerbation, and suicidal behavior in patients with panic disorder, 103-105
Present State Examination, assessment of negative symptoms in schizophrenia, 192-193
Principal component analysis, determination of covariation of DMS-IV personality disorders in mixed psychiatric sample, 206-215
PSE, *see* Present State Examination
Pseudoakathisia
case reports, 70-72
definition, 70
PSG, *see* Polysomnography
PSNAS, *see* Personal Social Network Adjustment Scale
PSS-Fam/Fr, *see* Perceived Social Support Family/Friends Scale
PSS-I, *see* PTSD Symptom Scale Interview
PSUD, *see* Psychoactive substance use disorder
Psychiatric diagnosis
associated with irritable bowel syndrome, 119
comorbidities in alcoholic patients, 334-338
Psychiatric Diagnostic Interview, assessment of comorbid psychiatric diagnosis and long-term drinking outcome, 335
Psychiatric disorders, associated with PTSD, 469-478
Psychiatric states, computerized measurement, 326-333
Psychoactive substance use disorder, comorbidity in eating disorder patients, 454-460
Psychoevolutionary theory of emotions, 35
Psychopathology
following civilian trauma, 20
relationship to alcohol intake and long-term outcome, 337
in Vietnam War veterans, 22
Psychosis, therapeutic nature of delirium, 450-453
Psychotic episodes, nature of in borderline personality disorders, 43
Psychotic phenomena, self-reported, associated with personality disorders, 42-48
PTCA, *see* Parent-to-child aggression
PTSD, *see* Posttraumatic stress disorder
PTSD Symptom Scale Interview, assessment of PRSD symptoms in sexual assault survivors, 50
Purposeful action, impaired in first-episode schizophrenia, 385-391
Pyschiatric patients, irritable bowel syndrome in, 116-122

Rape
association with anxiety disorder, 317
MDD, GAD, and substance dependence following, 21
see also Sexual assault
Rapid eye movement sleep, phasic, disturbance in patients with PTSD, 469-478
Rathus Assertiveness Scale, assessment of personality disorders in patients with body dysmorphic disorder, 230-231
Rearing style, affectionless control, and drug-dependent patients, 123-129

Reboxetine
assessing efficacy of in improving social function in patients with depression, 63-69
compared with fluoxetine in improving social functioning, 66
REHAB, *see* Rehabilitation Evaluation Hall and Baker
Rehabilitation Evaluation Hall and Baker, assessment of negative symptoms in schizophrenia, 192-193
Reitan Neuropsychological Assessment Metric Battery, scores for cognitive impairment compared with computer measurement scores, 326-333
REM sleep, *see* Rapid eye movement
Reward dependence, TPQ dimensions in adolescents, 426-431
Ritualistic behavior, in trichotillomania, 350
Rorschach variables, and performance of first-episode schizophrenic patients on WCST, 388

SAD, *see* Seasonal affective disorder
SADS, *see* Schedule for Affective Disorders and Schizophrenia
SANS, *see* Scale for the Assessment of Negative Symptoms
SASS, *see* Social Adaptation Self-evaluation Scale
SASS, *see* Social Adaptation Self-evaluation Scale
SBD, *see* Sleep breathing disorders
Scale for the Assessment of Negative Symptoms, in schizophrenia, 192-193
Scale to Assess Unawareness of Mental Disorders, assessment of
insight and outcome in bipolar, unipolar, and anxiety disorders, 167
psychotic biopolar disorder, 243
SCALUP, axis I diagnoses, 316
SCAN, *see* Schedule for Clinical Assessment in Neuropsychiatry
Schedule for Affective Disorders and Schizophrenia
Lifetime Version, assessment of comorbidity in bipolar II disorder, 339-343
suicide subscale, assessment of suicidal behavior in patients with panic disorder, 103
Schedule for Clinical Assessment in Neuropsychiatry, assessment of basic symptoms of schizophrenia, 285-286
Schizoid personality disorder
concordance rate in twin studies, 418
criteria for, 139
Schizoid personality trait, TPQ dimensions in adolescents, 428
Schizoid traits, associated with BDD, 5
Schizophrenia
assessment of
defense mechanisms in adolescents with, 35-41
treatment relationship, 197-205
association between dermatoglyphic ridge counts and cerebral structural abnormalities, 380-384
basic symptoms, effect of neuroleptic dosage and extrapyrimidal side effects, 284-288
and catatonia, 147
characteristics of patient and treatment relationship, 202
discrepancy between examiner rating and self-rating of negative symptoms, 191-916
expressed emotion and family distress in relatives of patients with, 392-397
extrapyramidal symptoms and neuroleptic treatment, 284-288
first-episode, relationship between duration of untreated psychosis and prognosis, 184-190
Schizophrenia (*Continued*)
patient perceptions of therapist, 204
predictors of discrepancy between self-rating and examiner rating for negative symptoms, 194
see also Negative symptoms
subjective experience and response to neuroleptics, 446-449
temporal sequence of psychotic symptoms, 187, 188-189
therapist perceptions of patient, 204
Wisconsin Card Sorting Test performance impaired in, nature of, 385-391
Schizotypal personality disorder
associated with history of trauma, 321
criteria for, 139
in male obsessive-compulsive disorder patients, 266-272
symptoms, self-reported, correlations with DSM-III-R personality disorders, 46
Schizotypal personality trait, TPQ dimensions in adolescents, 428-429
SCID II, *see* Structured Clinical Interview for DSM
SD, *see* Semantic differential
Seasonal affective disorder
comparison of rates for Western countries and China, 57
evidence for two forms of syndrome, 61
Seasonal Pattern Assessment Questionnaire, survey of seasonal changes in mood and emotion in Chinese college students, 57-62
Seasonality patterns
in Chinese college students, 57-62
winter and summer, characteristics, 58
Self-defeating personality disorder
associated with PTSD, 315-325
diagnosis, 321-323
concordance rate in twin studies, 418
criteria for, 139
Self-defeating personality trait, TPQ dimensions in adolescents, 430
Self-disclosure, in trichotillomania female patients, 349
Self-mutilative behavior, in dissociative disorders, psychiatric outpatients in Turkey, 216-222
Semantic differential technique, assessment of treatment relationship in schizophrenia, 197-205
Sensitivity, psychological symptoms during adjustment in immigrants, 291, 292
Separation anxiety, excessive, risk factor for personality disorders in panic disorder patients, 28
Serotonergic function, in social anxiety disorder, 408
Serotonin receptors, postsynaptic, dysfunction in social anxiety disorder, 405-415
Serotonin-selective reuptake inhibitors, treatment, social anxiety disorder, 408, 409, 410
Sertraline, treatment, social anxiety disorder, 410
Sexual abuse, childhood
definition, 131
personality traits of women with history of, 130-136
Sexual-assault survivors, PTSD and sleep breathing and movement disorders, 49-56
Shame, in trichotillomania female patients, 346
Simple phobia, in major depressive disorder, 97-102
Sleep
breathing disorders, assessment of in sexual assault survivors, 51

Sleep (*Continued*)
breathing and movement disorders in sexual assault survivors, 49-56
decreased need for, in mixed versus pure mania, 239
disorders, in PTSD, 475-476
disturbances, associated with PTSD, 469-478
movement disorders, assessment of in sexual assault survivors, 51
paralysis, in PTSD, 473
talking, in PTSD, 473
Sleep-Eval system, assessment of sleep disorders, 470
SMD, *see* Sleep movement disorders
Social Adaptation Self-evaluation Scale
as outcome measure in depression, 66-68
assessment of social functioning, 65
self-assessment of social functioning by patients with depression, 63-69
Social anxiety disorder
comorbidity, 407-408
social phobia, epidemiology, diagnosis, neurobiology, and treatment, 405-415
summary of SSRI studies, 411
treatment options, 409, 411-412
Social factors, differences in cluster B personality disorders, 435
Social functioning
assessment of in depression, 63-69
gender-related impairment in OCD, 269
methods of assessment, 64-65
and recovery from depression, 63-64
Social phobia
associated with BDD, 4
comorbidity in eating disorder patients, 454-460
in major depressive disorder, 97-102
see also Taijin Kyofusho
and social anxiety disorder, epidemiology, diagnosis, neurobiology, and treatment, 405-415
Social support, associated with major depression, 93
Social withdrawal, and duration of untreated psychosis in first-episode schizophrenia, 186
Somatization, predictor of remission in DSM hypochondriasis, 179
Somatoform disorder, *see* Body dysmorphic disorder
SPAQ, *see* Seasonal Pattern Assessment Questionnaire
Speilberg's State-Trait Anxiety Inventory, screening for binge eating disorder in obese outpatients, 111
SSIAM, *see* Structured and Scaled Interview to Assess Maladjustment
SSRI, *see* Serotonin-selective reuptake inhibitors
STAI, *see* Speilberg's State-Trait Anxiety Inventory
STAI, *see* State-Trait Anxiety Inventory
State-Trait Anxiety Inventory, assessment of anxiety and depression in OCD, 267
Stress
chronic, and major depression, 95
see also Civilian trauma; Posttraumatic stress disorder
Striatal presynaptic dopamine transporter, dysfunction in social anxiety disorder, 405-415
Stroop Color and Word Test, scores for cognitive impairment compared with other instruments, 326-333
Structured and Scaled Interview to Assess Maladjustment, assessment of social functioning, 64-65
Structured Clinical Interview for DSM
diagnoses and personality impairment scores in BDD patients, 8
evaluation of personality impairment in BDD patients, 4-12
Structural Clinical Interview for DSM-III-R
assessment of
depersonalization in panic disorder, 172-173
personality disorders, 137
psychotic bipolar disorder, 243
substance abuse, 277
Axis I, assessment of comorbidity in bipolar II disorder, 339-343
Patient Edition, assessment of comorbid anxiety disorders in MDD, 98
Personality Disorders
assessment
in conjunction with personality and substance abuse disorders, 277
in OCD, 267
personality disorder in patients with body
dysmorphic disorder, 230-231
relationship with TPQ, personality traits in adolescents, 426-431
twin studies using, 415-425
Structured Clinical Interview for DSM-IV
Axis Disorders, diagnosis of binge eating disorders, 160
Axis I Disorders, use in distinguishing borderline personality disorder from bipolar II disorder, 107
Axis II Personality Disorders
diagnosis of personality disorders in PDAG, 28-34
use in distinguishing borderline personality disorder from bipolar II disorder, 107
Structured Interview for Schizotypy, assessment of early- and late-onset OCD, 374
Substance abuse
comorbidity with social anxiety disorder, 407-408
following civilian trauma, 21
Substance use disorder, predictor of criminality, 276-284
Suicide
attempts
associated with BDD, 4
correlation with polarity of first episode in bipolar I disorder, 13-18
in dissociative disorders, psychiatric outpatients in Turkey, 216-222
and premenstrual exacerbation in patients with panic disorder, 103-105
see also Parasuicide
SUMD, *see* Scale to Assess Unawareness of Mental Disorders
Summer, rates of SAD during in Chinese college students, 57-62
Substance abuse, in Vietnam War veterans, 22
Symbol Digit Modalities Test, scores for cognitive impairment compared with other instruments, 326-333
Symptom Checklist 90, assessment of comorbid psychiatric diagnosis and long-term drinking outcome, 335
Symptom Questionnaire, assessment of comorbid anxiety disorders in MDD, 98

Taijin Kyofusho, culturally distinctive phobia in Japan, 457
Talbieh Brief Distress Inventory, assessment of psychological symptoms during adjustment in immigrants, 289-290

TAS-20, *see* Toronto Alexithymia Scale-20
TBDI, *see* Talbieh Brief Distress Inventory
Temporal lobe, implication in pathogenesis of panic disorder, 176
The American Psychopathological Association, program of 90th annual meeting, 1-3
Therapists, assessment of treatment relationship in schizophrenia, 197-205
Thyroid function, in mixed mania, 85
Toronto Alexithymia Scale-20, assessment of hemispheric functioning and gender, 352-359
Torture, PTSD, anxiety, and depression following, 20-21
Toxic chemical exposure, depression, alcohol dependence, phobia, GAD, and PTSD following, 21
Transsexualism, *see* Gender identity disorder
Trauma
  association with anxiety disorders, 315-325
  childhood, relationship to adult personality disturbance, 29
  civilian
    compared with combat, 19-20
    psychiatric sequelae, 19-23
  combat-related, personality disorders associated with, 322-323
  demographic and clinical characteristics, 318
  see also Posttraumatic stress disorder
Traumatic event
  association with anxiety disorder, 317
  definition of, 475
Traumatic experiences, associated with major depression, 93
Traumatic personality, in Vietnam combat veterans, 315
Trichotillomania
  psychosocial issue for in women, 344-351
  triggering events for, 349-350
Tricyclic agents, treatment, social anxiety disorder, 409
Tridimensional Personality Questionnaire, and relationship with DSM-III-R, personality traits in adolescents, 426-431
Twin studies, personality disorders, 416-425

Unipolar disorder
  insight and outcome in, 167-177
  major depressive episode, comparison with bipolar II patients, 107
Unipolar major depressive disorder, insight and outcome in, 169

Valproate, exposure to in bipolar disorder outpatients, 249-250
Venlafaxine, treatment, social anxiety disorder, 412
Vietnam War veterans, EMDR therapy for combat-related PTSD, 24-27

WAIS-R, performance of first-episode schizophrenic patients, 387
WCST, *see* Wisconsin Card Sorting Test
Winter, rates of SAD during in Chinese college students, 57-62
Wisconsin Card Sorting Test, impaired performance in first-episode schizophrenia, nature of, 385-391
Women
  alexithymia and hemispheric functioning in, 356-357
  diagnosis of mixed mania in, 83-85
  premenstrual exacerbation of suicidal behavior in patients with panic disorder, 103-105
  psychosocial issues for trichotillomania in, 344-351
  see also Gender

Yale Tic Inventory, assessment of early- and late-onset OCD, 374
Yale-Brown Obsessive-Compulsive Scale, assessment of
  early- and late-onset OCD, 374
  OCD, 267
Y-BOCS, *see* Yale-Brown Obsessive-Compulsive Scale

Zung's Self-Rating Depression Scale, assessment of anxiety and depression in OCD, 267